

Scale invariance in percolation theory and fractals.

*M.V. Entin, G.M. Entin.*Institute of Semiconductor Physics, Siberian Branch of the Russian Academy of Sciences,
630090 Novosibirsk, Russia.

Novosibirsk State University, 630090 Novosibirsk, Russia

E-mail:Entin@isp.nsc.ru

PACS 72.60.+g

The properties of the similarity transformation in percolation theory in the complex plane of the percolation probability are studied. It is shown that the percolation problem on a two-dimensional square lattice reduces to the Mandelbrot transformation, leading to a fractal behavior of the percolation probability in the complex plane. The hierarchical chains of impedances, reducing to a nonlinear mapping of the impedance space onto itself, are studied. An infinite continuation of the procedure leads to a fixed point. It is shown that the number of steps required to reach a neighborhood of this point has a fractal distribution.

In percolation theory, an approach based on a hierarchical chain of resistances is used to calculate the percolation threshold and the behavior of a system near the percolation threshold. The approach is based on a recurrence relation for the percolation probability p [1] of a medium or for the resistances of the hierarchically constructed cluster (see, for example, Ref. 2).

A very simple model of a scale transformation for geometric percolation employs a calculation of the percolation probability for a d -dimensional cube with edge length $2a$ and known percolation probability a . If one neglects percolation between the diagonally positioned conducting cubes, the renormalization-group relation for the percolation probability at the n th level has the form [1]:

$$p_n = 1 - (1 - p_{n-1}^2)^{2^{d-1}}. \quad (1)$$

The mapping (1) is a particular case of more general formulations of the renormalization-group study of percolation transitions. Our objective in the present letter is to investigate the mapping (1) in the complex p -plane. This mapping is nonlinear and its general properties are described by the theory of fractals [3]. Another problem is to study a similar mapping for hierarchically constructed chains of impedances simulating the high-frequency properties of a percolation medium.

Let us consider the two-dimensional (2D) case. The mapping (1) possesses three fixed points: 0, 1, and $p = (\sqrt{5} - 1)/2$. The point p_c is unstable: for $p_0 > p_c$, p_n asymptotically approaches 1, and in the opposite case p_n approaches zero. Therefore p_c is the percolation threshold. The investigation of fractals is based on the study of the very simple nonlinear Mandelbrot map

$$x' = F(x) = x^2 + c. \quad (2)$$

It is easy to see that choosing $c = -1$, substituting Eq. (2) into itself, and replacing x by $-p$ gives precisely the map (1) with $d = 2$. Therefore the Sarychev map (1) is a particular case of the Mandelbrot map. By studying this map in the complex p -plane we can determine the regions from which a conducting ($p_n \rightarrow 1$) or nonconducting ($p_n \rightarrow 0$) medium is reached as $n \rightarrow \infty$. In terms of the Mandelbrot theory, they are called basins of attraction of the corresponding points

The attraction basin of the point zero, corresponding to nonpercolation in the 2D case, is displayed on Fig. 1. It is distinguished by coloring the plane as a function of the parity of number of steps required to reach a circle with radius 310^{-2} around the origin. The basins of attraction of other finite fixed points are colored in black. The basin of attraction for zero cuts off $(0, p_c)$ of the real axis the part from the segment $(0, 1)$ corresponding to the physical region. The percolation system is determined by the segment $(p_c, 1)$, which falls in the basin of attraction of the point 1 - the black region on the right-hand side.

The fractal structure of the basin of attraction is formed by regions spanning the points that map into zero after many repetitions of the mapping, i.e. , the points are zeros of the functions $F_n(p) = F(F(\dots F(p))\dots)$. This function determines the percolation probability of a square with an edge length 2^n . The derivative of $F_n(p)$ with respect to p can be regarded as the distribution function of the percolation thresholds. F_n is a polynomial of degree 2^n with even powers.

It is easy to obtain the asymptotic behavior of F_n as $\delta p = p - p_c \rightarrow 0$. Linearizing the map (2) gives $\delta p' = 2p_c \delta p$ i.e. $F'_n(p_c) \simeq (2p_c)^n$. Since $2p_c > 1$, the derivative grows exponentially with n and $F_n(p) \rightarrow \Theta(p - p_c)$ as $n \rightarrow \infty$.

The general form of this function is, however, quite complicated. It can be shown that the general formula for the zeros of this polynomial has the form

$$\sigma_1 \sqrt{1 + \sigma_2 \sqrt{1 + \sigma_3 \sqrt{\dots + \sigma_n}}} \quad (3)$$

where $\sigma_i = \pm 1$. Evidently, all zeros before the mappings are also zeros after the mappings. The preceding zeros appear with a higher multiplicity. Specifically, if $n = 2^k$, then the zero of the 2^j th mapping ($j < k$) appears with multiplicity 2^{k-j} .

In the limit $n \rightarrow \infty$ the zeros approach the boundary of the basin of attraction, forming a so-called Julia set. The fractal nature of the Julia set can be discovered by examining the periodic subsequences $\{\sigma_i\}$. Each such subsequence approaches its own irrational limit. Together the limits all form a fractal boundary.

It is easy to obtain a physical interpretation of a basin of attraction. Consider the quantity $Re(\ln F_n(x))$. This quantity is 2D electrostatic potential of charges which are placed at zeros of the function $F_n(x)$ and have a magnitude equal to the multiplicity of the zeros. The equipotential lines form the basin of attraction. Large, circular regions of influence of the charges arise near the largest charges-the zeros of the mappings with small j . As j increases, the number of regions increases as the number of zeros and size ρ_n of regions decreases. The rate of decrease of the size can be found by equating the potential produced by the charges at the points where the boundaries of the corresponding regions touch:

$$2^{-j} \ln \rho_{2^j} = 2^{-(j+1)} \ln \rho_{2^{-(j+1)}}$$

whence $\rho_n = \rho_0^n$. The choice of the set $\{\sigma_n\}$ equal to 1 gives, in accordance with the Lee-Yang theorem, a chain of zeros of a polynomial which approach p_c .

Therefore the percolation probability behaves very simply on the axis in the limit of large n , viz. as $\Theta(p - p_c)$, and in the complex plane it exhibits a complicated, fractal behavior. If one is interested only the asymptotic behavior of $F_n(p)$ for large n and near a transition, then there is no need to know the singularities of this function in the complex plane. However, if we wish to describe the function for all n and far from a transition point, then knowing the zeros of $F_n(p)$ makes it possible to reconstruct the entire function.

Another example of the fractal behavior is the frequency dependence of the impedance of a large, hierarchically constructed circuit in the case of small, real losses. In Ref. 4 it was found that in a random medium consisting of capacitors, inductors and resistors with small resistance, the local electric field is subject to anomalously strong fluctuations.

For example, in a 2D Dykhne medium consisting of two metals the imaginary part of the effective permittivity at a frequency between their plasma frequencies is nonzero in the absence of local absorption. This causes the local electric field to diverge. We shall show that anomalously strong fluctuations in the properties occur even in a regular hierarchical chain. Strictly speaking in the absence of absorption even self-averaging of the system does not occur.

Consider now the electrical circuits in Fig.2. Let the initial element i be a capacitor C with impedance $Z_i = i/\omega C$ at frequency ω and the impedance of the initial element of the iteration sequence $Z_0 = -iL\omega + R$, where L is the inductance and R is the resistance. Switching to dimensionless variables $z_n = \omega C Z_n$, $z_0 = -iLC\omega^2 + RC\omega$, we obtain for the circuit in Fig.2a

$$z_{n+1} = i \frac{(2z_n + i)}{(z_n + i)}. \quad (4)$$

The explicit formula for z_n has the form $z_n = (if_{n+1}z_0 - f_n)/(f_n z_0 + if_{n-1})$, where f_n are the Fibonacci numbers

$$f_n = \frac{1}{\sqrt{5}} \left[\left(\frac{1 + \sqrt{5}}{2} \right)^{n+1} - \left(\frac{1 - \sqrt{5}}{2} \right)^{n+1} \right]$$

The set of both poles and zeros has a limit point $-i(\sqrt{5} - 1)/2$. Near this point z_n converges poorly.

In the example analyzed above, the final formula reduces to a linear fractional transformation. In the general case the map (2) performed n times leads to rational function of high order. The result of this mapping in the space z_0 can be predicted if the stationary points and poles of transformation are known. As a degree of the transformation increase, the number of poles increase as 2^n and the convergence of the mapping therefore becomes a fractal function of the initial impedance.

Fig.2b corresponds to the map

$$z_{n+1} = \frac{z_n(z_n + i)}{2z_n + i}. \quad (5)$$

Figure 3 displays the impedance after 10 iterations as a function of z_0 . It is evident from the figure that in the absence of absorption the impedance is a fractal curve. According to the definition of z_0 , the fractal nature is reproduced in the frequency dependence of the impedance. The fractal behavior is explained by the fact that random circuits which resonate at the frequency ω arise in the system. In the limit of small R , the width of a resonance is determined by the quantity R/L . The total number of resonances distributed in a characteristic frequency band of the order of $\sqrt{LC}$ is determined by the number $2^n + 1$ of the element in the system. As the static conductivity of elements increases, the resonances cease to overlap if $\sqrt{LC}2^{-n}L/R > 1$, and the absorption is smoothed out.

An important property of a percolation system is the percolation radius r_c . The geometric percolation radius is determined as the scale on which the probability of two sites belonging to the same cluster starts to decrease more rapidly than in power law. The correlation radius becomes infinite at percolation threshold.

In the analysis of the conductivity of a two-phase random system containing phases with finite dc conductivities, it is assumed that the correlation radius is related to the ratio $h = \sigma_1/\sigma_2$ of the conductivities: $r_c \sim h^{-\beta}$. However, the question of what determines the correlation radius remains open in the case of an ac field. A necessarily real quantity, which r_c is, cannot be an

analytic function of the impedances. Therefore the function $r_c(\sigma_1, \sigma_2)$ cannot be suggested as an alternative.

In the model corresponding to Fig. 2b, the impedance approaches zero as the number of iterations increases. In the limit of a sufficiently large number of steps, the renormalization-group equation can be converted into the differential equation $dz/dn = -z^2$ with the solution $z = 1/(n + A)$. It is natural to take as the correlation length, in units of n , the value of n for which z reaches this asymptotic law. For real z , this occurs for $n \sim \ln z$.

In Fig. 4, the number of iterations at which the impedance drops below 0.01 in modulus is displayed as a function of the initial impedance. One can see how this dependence becomes fractal as R decreases.

The correlation radius estimated from the smoothing of the resonances has the form $n \sim \ln(1/R)/\ln 2$. The fact that r_c becomes infinite can be understood if the system under study is interpreted in the limit $R \rightarrow 0$ as a system of coupled, localized states for a photon. In such a formulation, the spectrum of the system consists of separate lines, which do not vanish in the absence of a relaxational interruption of the phases. At the same time, the substitution $z \rightarrow iz$ gives the system coupled positive and negative resistances, i.e., the system contains amplifying sections, which explains the absence of a thermodynamic limit.

This work was partly supported by the Russian Fund for Fundamental Research, Grants Nos. 95-02-04432 and 96-02-19353 and the Volkswafen Foundation.

- [1] A. K. Sarychev, Zh. Eksp. Teor. Fiz. **72**, 1001 [Sov. Phys. JETP **45**, 524 (1977)].
- [2] J. P. Clerc, G. Giraud, J. M. Laugier and J. M. Luck, Adv. Phys. **39**, 191 (1990).
- [3] H. O. Peitgen and P. H. Richter, *The Beauty of Fractals: Images of Complex Dynamic Systems* Springer-Verlag, New York, 1986 [Russian Translation, Mir, Moscow, 1993, p. 176].
- [4] F. Brouers, S. Blacher, N. Henriouille, A. Sarychev, in *Electrical Transport and Optical Properties of inhomogeneous media*, Scientific Center for Applied Problems in Electrodynamics, Moscow, 1996, p. 46.
- [5] A. M. Dykhne, Zh. Eksp. Teor. Fiz. **59**, 110 (1970) [Sov. Phys. JETP **32**, 63 (1971)].

Picture signs

FIG. 1. a-Basin of attraction of the fixed point 0 corresponding to nonpercolation with $d = 2$. The regions where the number of steps required to reach a neighborhood of the point 0 is even are colored in black, as are the basins of attraction of other, nonfixed points, except infinity. b-Same as Fig. 1a, $d = 3$.

FIG. 2. Circuits of the hierarchical chains a and b corresponding to the expressions (4) and (5).

FIG. 3. Impedance of the chain in Fig. 2b as a function of the imaginary part of the initial impedance after 10 iterations. Adjacent curves, except for the bottom curve, are raised by 0.2 units relative to one another for better resolution.

FIG. 4. a-Number of steps required to reach a neighborhood of 310^{-2} as a function of the initial impedance for the circuit in Fig. 2b. Adjacent curves, except for the bottom curve, are raised by 25 units relative to one another. b-Dual coloring, similar to Fig. 1, of the plane of initial impedances.




Fig. 1




Fig. 2




Fig. 3




Fig. 4a



Fig. 4b